

---

---

## Worksheet 1: Fundamentals

### Objectives

- Convert between units.
- Carry out mathematical operations with vectors.

### Summary

#### *Quantities and units*

All the phenomena we will discuss in this class can be described in terms of three fundamental quantities: **distance**, **mass**, and **time**. The respective mks or SI units for these are the **meter** m, **kilogram** kg, and **second** s. The units for all physical quantities used in this course can be expressed in terms of these fundamental units.

Units of measurement can be operated on mathematically just like variables in algebra—they add together, subtract from each other, multiply, and divide. Converting between units of the same quantity requires identifying a proportional relationship between them.

Example: 1 h = 60 min, so 24 h = 24 (60 min) = 1440 min.

#### *Vectors*

Many physical quantities have particular directions and are expressed as **vectors**.

Often expressed as **components**:  $\vec{A} = (A_x, A_y, A_z) = A_x\hat{i} + A_y\hat{j} + A_z\hat{k}$ .

**Unit vectors**:  $\hat{i} = (1, 0, 0)$ ;  $\hat{j} = (0, 1, 0)$ ;  $\hat{k} = (0, 0, 1)$

**Magnitude**:  $\|\vec{A}\| = A = \sqrt{A_x^2 + A_y^2 + A_z^2}$

**Addition**:  $\vec{A} + \vec{B} = (A_x + B_x, A_y + B_y, A_z + B_z)$

**Scalar multiplication**:  $c\vec{A} = c(A_x, A_y, A_z) = (cA_x, cA_y, cA_z)$

**Scalar product (dot product)**:  $\vec{A} \cdot \vec{B} = A_xB_x + A_yB_y + A_zB_z = AB\cos(\theta)$ , where  $\theta$  is the angle from  $\vec{A}$  to  $\vec{B}$ .

**Vector product (cross product)**:  $\vec{A} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ A_x & A_y & A_z \\ B_x & B_y & B_z \end{vmatrix}$ ;  $\|\vec{A} \times \vec{B}\| = AB\sin(\theta)$ .

$\vec{A} \times \vec{B}$  is a vector, perpendicular to both  $\vec{A}$  and  $\vec{B}$ , in the direction given by the right-hand rule.

## Problems

1. What are the units of volume? Suppose another student tells you that a cylinder of radius  $r$  and height  $h$  has a volume given by  $\pi^3 h$ . Explain why this cannot be right.
2. Light travels in a vacuum  $2.9979 \times 10^8$  m in 1 s. How many nanoseconds does it take to travel 1.00 ft?
3. If the scalar product of two vectors is negative ( $\vec{A} \cdot \vec{B} < 0$ ), what does that tell you about them?
4. The magnitude of the vector product of two vectors is proportional to the sine of the angle between them. Vector magnitudes are always considered positive, yet a sine can be negative as well as positive. If  $\sin(\theta)$  of the angle from  $\vec{A}$  to  $\vec{B}$  is negative:
  - a. What does that mean about the angle  $\theta$ ?
  - b. What does that mean for the vector  $\vec{A} \times \vec{B}$ ?
5. Define  $\vec{L}$  to be the vector running along the length of this room,  $\vec{W}$  to be the room's width vector, and  $\vec{H}$  to be its height vector.
  - a. What is the approximate value of  $\vec{L} \times \vec{W}$ , the vector product of the length and width vectors?
  - b. What is the physical significance of  $\vec{L} \times \vec{W}$ ?
  - c. What is the approximate value of  $(\vec{L} \times \vec{W}) \cdot \vec{H}$ ?
  - d. What is the physical significance of  $(\vec{L} \times \vec{W}) \cdot \vec{H}$ ?